

Low scale Seesaw model and Lepton Flavor Violating Rare B Decays

T. Fujihara^(a) *, S. K. Kang^(b) †, C. S. Kim^(a,c) ‡, D. Kimura^(a) § and T. Morozumi^(a) ¶

(a): *Graduate School of Science, Hiroshima University,*

Higashi-Hiroshima, Japan, 739-8526

(b): *Seoul National University, Seoul, Korea*

(c): *Department of Physics, Yonsei University, Seoul 120-749, Korea*

Abstract

We study lepton flavor number violating rare B decays, $b \rightarrow sl_h^\pm l_l^\mp$, in a seesaw model with low scale singlet Majorana neutrinos motivated by the resonant leptogenesis scenario. The branching ratios of inclusive decays $b \rightarrow sl_h^\pm \bar{l}_l^\mp$ with two almost degenerate singlet neutrinos at TeV scale are investigated in detail. We find that there exists a class of seesaw model in which the branching fractions of $b \rightarrow s\tau\mu$ and $\tau \rightarrow \mu\gamma$ can be as large as 10^{-10} and 10^{-9} within the reach of Super B factories, respectively, without being in conflict with neutrino mixings and mass squared difference of neutrinos from neutrino data, invisible decay width of Z and the present limit of $Br(\mu \rightarrow e\gamma)$.

* fujihara@theo.phys.sci.hiroshima-u.ac.jp

† skkang@phya.snu.ac.kr

‡ cskim@yonsei.ac.kr

§ kimura@theo.phys.sci.hiroshima-u.ac.jp

¶ morozumi@hiroshima-u.ac.jp

I. INTRODUCTION

Thanks to the worldwide collaborative endeavor for B factories and neutrino experiments, our understanding of the flavor mixing phenomena in quark sector as well as in lepton sector has been dramatically improved over the past few years. Even though discoveries of neutrino oscillations in solar, atmospheric and reactor experiments gave a robust evidence for the existence of non-zero neutrino masses, we do not yet understand mechanisms of how to generate the masses of neutrinos and why those masses are so small. The most attractive proposal to explain the smallness of neutrino masses is the seesaw mechanism [1, 2, 3, 4] in which super-heavy singlet particles are introduced. One of the virtues of the seesaw mechanism is to provide us with an elegant way to achieve the observed baryon asymmetry in our universe via the related leptogenesis [5]. However, the typical seesaw scale is of order $10^{10} \sim 10^{14}$ GeV, which makes it impossible to probe the seesaw mechanism at collider experiments in a foreseeable future. Moreover, the leptogenesis at such a high energy scale meets a serious problem, the so-called gravitino problem, when it realizes in supersymmetric extensions of the standard model. Thus, it may be quite desirable to achieve the seesaw mechanism as well as the leptogenesis at a rather low energy scale. In these regards, scenarios of the resonant leptogenesis, in which singlet neutrinos with masses of order $1 \sim 10$ TeV are introduced, have been recently proposed [6, 7, 8, 9, 10, 11, 12, 13]. Interestingly enough, the amplitudes of some flavor violating processes, which are highly suppressed in usual seesaw models with super-heavy singlet neutrinos, may be enhanced with such low scale singlet neutrinos. Thus, it is worthwhile to examine how we can probe a seesaw model with low scale singlet neutrinos at collider experiments and to find some experimental evidence for the seesaw model via probe of lepton flavor violating processes.

In this paper, within the context of a low scale seesaw model based on $SU(2)_L \times U(1)$ [14], we study quark and lepton flavor violating (QLFV) rare decay processes such as $b \rightarrow sl_h^\pm l_l^\mp$ and $l_h \rightarrow l_l \gamma$, where l_h and l_l denote heavy and light charged leptons, respectively. After the quark flavor changing neutral currents (FCNC) $b \rightarrow sl^+l^-$ had been discovered in B factories [15], it has been naturally expected that the next generation experiments could probe well the lepton flavor violating (LFV) processes [1, 16, 17, 18], which may eventually uncover the mechanism for the generation of small neutrino masses and the leptogenesis as well. Therefore, the precise predictions for those processes based on well motivated scenarios are very useful to find out if such scenarios can describe the nature correctly or not. Here we focus on the seesaw model motivated by the resonant leptogenesis scenario with two

almost degenerate singlet neutrinos at TeV scale. In the context of the seesaw mechanism, lowering the singlet mass scale leads to an undesirable enhancement of the light neutrino masses unless the Dirac neutrino couplings are guaranteed to be naturally small. However, as shown in [19, 20], despite of the low mass scale of singlet Majorana neutrinos we can obtain light neutrino mass spectrum consistent with the current neutrino data by tuning the phase of the Yukawa-Dirac mass terms so that the two degenerate singlets contribute to the low energy effective Majorana mass terms destructively and the lepton number is approximately conserved. Interestingly enough, in such a scenario the sizable LFV processes and the suppression of the lepton number violation required in the effective Majorana mass for light neutrinos may coexist. We will show that the amplitudes of quark FCNC and LFV processes can be enhanced by considering some specific structure of Yukawa-Dirac and singlet Majorana mass matrices which are required to achieve the resonant leptogenesis. We will obtain rather stringent constraints on QLFV processes by taking the constraints arisen from the invisible decay of Z boson, neutrino mass-squared differences and lepton flavor mixings measured at neutrino experiments and the experimental constraints of LFV processes such as $l_h \rightarrow l_l \gamma$.

The paper is organized as follows: In section 2, we present the lepton flavor mixings of seesaw models with arbitrary number of singlet Majorana neutrinos. The analytical expressions for the branching fractions of the QLFV rare B decays are presented and the model independent bound on the branching fractions is obtained. In section 3, we study the low mass scale scenarios for singlet neutrinos and build the models in which the quark FCNC and LFV processes are enhanced. We give the predictions for the QLFV processes by taking into account the various constraints. In section 4, we summarize the results.

II. LEPTON FLAVOR MIXING OF SEESAW MODEL AND QLFV RARE B DECAY

The lepton flavor mixings of seesaw model are described in detail in Refs. [21, 22]. Here we extend the model to the case with arbitrary number of singlet neutrinos and introduce a convenient decomposition of the Yukawa-Dirac mass term which is useful to our study. Let us start with a seesaw model described by following Lagrangian,

$$\mathcal{L}_m = -y_\nu^{ik} \overline{L}_i N_{R_k} \tilde{\phi} - y_l^i \overline{L}_i l_{R_i} \phi - \frac{1}{2} \overline{N_{R_k}^c} M_k N_{R_k} + h.c., \quad (1)$$

and the neutrino mass matrix

$$M_\nu = \begin{pmatrix} 0 & m_D \\ m_D^T & M \end{pmatrix}, \quad (2)$$

where M is a $N \times N$ real diagonal singlet Majorana neutrino mass matrix and m_D is a Dirac Yukawa mass term. Then, $(3 + N) \times (3 + N)$ neutrino mass matrix can be diagonalized by the mixing matrix $\mathcal{V}$ as,

$$M_\nu^{\text{Diag}} = \mathcal{V}^\dagger M_\nu \mathcal{V}^*. \quad (3)$$

If the seesaw condition, *i.e.* $\frac{m_D}{M} < 1$, is satisfied, $(3 + N) \times (3 + N)$ unitary matrix $\mathcal{V}$ can be approximately parameterized as,

$$\mathcal{V} = \begin{pmatrix} V & m_D \frac{1}{M} \\ -\frac{1}{M} m_D^\dagger V & (1_N - \frac{1}{M} m_D^\dagger m_D \frac{1}{M})^{\frac{1}{2}} \end{pmatrix}, \quad (4)$$

where $\mathcal{V}$ satisfies unitarity to the order of $\frac{m_D^2}{M^2}$, and 1_N denotes an $N \times N$ unit matrix. Here 3×3 submatrix V is not exactly unitary and can be written [22] as

$$V = \left(1 - m_D \frac{1}{M^2} m_D^\dagger\right)^{\frac{1}{2}} V_0, \quad (5)$$

where V_0 is a unitary matrix which diagonalizes the effective Majorana mass term,

$$\begin{aligned} m_{eff} &= -m_D \frac{1}{M} m_D^T, \\ V_0^\dagger m_{eff} V_0^* &= \text{Diag}[n_1, n_2, n_3], \end{aligned} \quad (6)$$

where n_i are the masses of three light neutrinos.

For convenience, we introduce the following parametrization for m_D [23],

$$m_D = (\mathbf{m}_{D1}, \mathbf{m}_{D2}, \dots, \mathbf{m}_{DN}) = \begin{pmatrix} \mathbf{u}_1 & \mathbf{u}_2 & \dots & \mathbf{u}_N \end{pmatrix} \begin{pmatrix} m_{D1} & 0 & 0 & 0 \\ 0 & m_{D2} & 0 & 0 \\ 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & m_{DN} \end{pmatrix}, \quad (7)$$

where N unit vectors are introduced as,

$$\mathbf{u}_I = \frac{\mathbf{m}_{DI}}{m_{DI}}, \quad (8)$$

with $m_{DI} = |\mathbf{m}_{DI}|$. By introducing the parameters of mass dimension $X_I = \frac{m_{DI}^2}{M_I}$ ($I = 1 \sim N$), m_{eff} can be written as

$$m_{eff} = - \sum_{i=1}^N \mathbf{u}_i X_i \mathbf{u}_i^T. \quad (9)$$

The charged current is associated with $3 \times (3 + N)$ submatrix of $\mathcal{V}$. The deviation from the unitarity of 3×3 matrix V is given as

$$\begin{aligned} \sum_{a=1}^3 V_{ia} V_{ja}^* &= \delta_{ij} - \sum_{I=1}^N \frac{X_I}{M_I} u_{iI} u_{jI}^*, \\ \sum_{i=e,\mu,\tau} V_{ia} V_{ib}^* &= \delta_{ab} - \sum_{I=1}^N (V_0^\dagger U)_{bI} \frac{X_I}{M_I} (U^\dagger V_0)_{Ia}, \end{aligned} \quad (10)$$

where $U = (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_N)$.

Now we study the QLFV rare B decay processes $b \rightarrow s l_h^\pm l_l^\mp$ in the context of the seesaw model we consider. We denote l_h as a heavy lepton and l_l as a light lepton, and the possible combinations for $(l_h^\pm, l_l^\mp)$ are $(\tau^\pm, \mu^\mp)$, $(\mu^\pm, e^\mp)$ and $(\tau^\pm, e^\mp)$. By separating the contributions from the light neutrinos with masses n_a ($a = 1 \sim 3$) and ones from the heavy neutrinos with masses M_{A-3} ($A = 4, 5, \dots, 3 + N$), we can express the amplitude for $b \rightarrow s l_h^- l_l^+$ as

$$\begin{aligned} T &= \frac{\sqrt{2} G_F \alpha_{QED}}{4\pi s_W^2} \times \\ &(\overline{u}_s \gamma_\mu L u_b) (\overline{u}_h \gamma^\mu L v_l) \sum_{i=u \ c \ t} V_{ib} V_{is}^* \left(\sum_{a=1}^3 V_{ha} V_{la}^* E(x_i, y_a) + \sum_{A=4}^{(3+N)} \mathcal{V}_{hA} \mathcal{V}_{lA}^* E(x_i, y_{A-3}) \right), \end{aligned} \quad (11)$$

where u_b, u_s, u_h and v_l denote the spinor of bottom quark, strange quark, heavy lepton and light anti-lepton respectively and $L = \frac{1-\gamma_5}{2}$, $x_i = \frac{m_i^2}{M_W^2}$, $y_a = \frac{n_a^2}{M_W^2}$, $y_A = \frac{M_A^2}{M_W^2}$. E is a Inami-Lim [24] function presented as,

$$\begin{aligned} E(x, y) &= -(1 + \frac{xy}{4}) \frac{f(x) - f(y)}{x - y} + (1 - \frac{7xy}{4}) \frac{xf'(x) - yf'(y)}{x - y}, \\ \text{with } f(x) &= \frac{x \log[x]}{x - 1}. \end{aligned} \quad (12)$$

We may neglect the up quark loop contribution because of the smallness of $V_{ub} V_{us}^*$. By using the unitarity relations for leptonic and quark sectors,

$$\begin{aligned} \sum_{a=1}^3 V_{ha} V_{la}^* &= - \sum_{A=4}^{3+N} \mathcal{V}_{hA} \mathcal{V}_{lA}^*, \\ V_{tb} V_{ts}^* &= -V_{cb} V_{cs}^*, \end{aligned} \quad (13)$$

we can simplify Eq. (11) as follows;

$$T = \frac{\sqrt{2}G_F}{4\pi} \frac{\alpha_{QED}}{s_W^2} (\bar{u}_s \gamma_\mu L u_b) (\bar{u}_h \gamma^\mu L \nu_l) V_{tb} V_{ts}^* \sum_{A=4}^{3+N} \mathcal{V}_{hA} \mathcal{V}_{lA}^* (\bar{E}(x_t, x_c, y_{A-3})), \quad (14)$$

In Eq.(14), we only keep the contributions of the top quark, charm quark and heavy neutrinos, and

$$\begin{aligned} \bar{E}(x_t, x_c, y_{A-3}) &= E(x_t, y_{A-3}) - E(x_t, 0) - E(x_c, y_{A-3}) + E(x_c, 0) \\ &\simeq x_t y_{A-3} \left(\frac{3}{4(1-x_t)(1-y_{A-3})} - \frac{(x_t^2 - 8x_t + 4) \log(x_t)}{4(x_t - 1)^2(x_t - y_{A-3})} - \frac{(y_{A-3}^2 - 8y_{A-3} + 4) \log(y_{A-3})}{4(y_{A-3} - 1)^2(y_{A-3} - x_t)} \right), \end{aligned} \quad (15)$$

which agrees with the results in Refs. [14, 25]. The branching fraction of $b \rightarrow sl_h^- l_l^+$ can be easily obtained as

$$\text{Br}(b \rightarrow sl_h^- l_l^+) = \text{Br}(b \rightarrow c \bar{\nu}_e e) \frac{|V_{tb} V_{ts}|^2}{|V_{cb}|^2} \frac{P(\frac{m_h}{m_b})}{P(\frac{m_c}{m_b})} \left(\frac{\alpha_{QED}}{8\pi s_W^2} \right)^2 \left| \sum_{A=4}^{3+N} \mathcal{V}_{hA} \mathcal{V}_{lA}^* \bar{E}(x_t, x_c, y_{A-3}) \right|^2, \quad (16)$$

where the phase factor P is given by $P(x) = 1 - 8x^2 + 8x^6 - x^8 - 24x^4 \log(x)$ and m_h denotes the heavier lepton (l_h) mass. For numerical computation, we take $\text{Br}(b \rightarrow c \bar{\nu}_e e) = 0.107$, $m_b = 4.75$ (GeV), $m_c = 1.25$ (GeV), $\alpha_{QED} = \frac{1}{137}$, $m_\tau = 1.78$ (GeV), $m_\mu = 0.106$ (GeV), $m_W = 80.4$ (GeV). The branching fractions are then given as

$$\begin{aligned} \text{Br}(b \rightarrow s \tau^- e^+) &= 1.0 \times 10^{-7} |S(\tau, e)|^2, \\ \text{Br}(b \rightarrow s \tau^- \mu^+) &= 1.0 \times 10^{-7} |S(\tau, \mu)|^2, \\ \text{Br}(b \rightarrow s \mu^- e^+) &= 2.8 \times 10^{-7} |S(\mu, e)|^2, \end{aligned} \quad (17)$$

where $S(h, l)$ is the suppression factor defined by

$$\begin{aligned} S(h, l) &= \sum_{A=4}^{3+N} \mathcal{V}_{hA} \mathcal{V}_{lA}^* \bar{E}(x_t, x_c, y_{A-3}), \\ &= \sum_{I=1}^N \frac{X_I}{M_I} \mathbf{u}_{hI} \mathbf{u}_{lI}^* \bar{E}(x_t, x_c, y_I). \end{aligned} \quad (18)$$

As can be seen above, the QLFV processes depend on $\frac{X_I}{M_I} = \frac{m_{D_I}^2}{M_I^2}$, $\mathbf{u}_{hI} \mathbf{u}_{lI}^*$ and M_I in $\bar{E}$. If $\frac{X_I}{M_I}$ is not very small, there will be a chance to detect the QLFV processes at B factories near future. First, we can roughly estimate the branching fractions of the QLFV processes.

We notice that a model independent constraint on $\sum_{I=1}^N \frac{X_I}{M_I}$ can be obtained from invisible decay width of Z and charged currents lepton universality test [26, 27, 28] because $Z \rightarrow \nu_a \bar{\nu}_b$ coupling is suppressed compared with standard model prediction as,

$$\frac{g}{2 \cos \theta_W} \sum_{a,b=1}^3 Z_{ba} Z^\mu \bar{\nu}_b \gamma_\mu \frac{(1 - \gamma_5)}{2} \nu_a, \quad (19)$$

where Z_{ba} is related to the violation of the unitarity of V ,

$$Z_{ba} = \sum_{i=e\mu\tau} V_{ib}^* V_{ia} = \delta_{ab} - (V_0^\dagger U)_{bI} \frac{X_I}{M_I} (U^\dagger V_0)_{Ia}. \quad (20)$$

In the model under consideration, the effective number of light neutrinos N_ν is given by

$$N_\nu = \sum_{a,b=1}^3 |Z_{ba}|^2 = 3 - 2 \sum_{I=1}^N \frac{X_I}{M_I}. \quad (21)$$

From the experimental result [29], $N_\nu = 2.984 \pm 0.008$, we can obtain the following bound,

$$\sum_{I=1}^N \frac{X_I}{M_I} = 0.008 \pm 0.004. \quad (22)$$

If the bound is dominated by N degenerate singlet heavy neutrinos with $X_1 \sim X_2 \sim X_N$, $\frac{m_{PI}}{M_I} \simeq \frac{0.1}{\sqrt{N}}$, which is achieved in the case of low scale singlet neutrinos.

From the fact that the lepton flavor universality of charged current is generally violated in seesaw model, one can obtain experimental bounds on $\epsilon_i \equiv \sum_{I=1}^N |u_{iI}|^2 \frac{X_I}{M_I}$ ($i = e, \mu, \tau$) from the lepton universality test of charged current interactions. The deviation from the universality is expressed in terms of the flavor dependent coupling g_i defined as

$$\frac{g_i^2}{g^2} = \sum_{a=1}^3 |V_{ia}|^2 = 1 - \epsilon_i. \quad (23)$$

Using the definition of g_i above, for instance, the decay width of W into charged lepton l_i and neutrino is given by,

$$\begin{aligned} \sum_{a=1}^3 \Gamma[W \rightarrow l_i \bar{\nu}_a] &= \frac{g^2 M_W}{48\pi} \sum_{a=1}^3 |V_{ia}|^2 \left(1 - \frac{m_i^2}{M_W^2}\right)^2 \left(1 + \frac{m_i^2}{2M_W^2}\right), \\ &= \frac{g_i^2 M_W}{48\pi} \left(1 - \frac{m_i^2}{M_W^2}\right)^2 \left(1 + \frac{m_i^2}{2M_W^2}\right), \end{aligned} \quad (24)$$

where we ignore the neutrino masses. The experimental bounds on $\epsilon_\mu - \epsilon_e$ and $\epsilon_\tau - \epsilon_e$ was obtained from the ratios of the branching fractions of W decays and also from the ratios of




FIG. 1: The experimentally allowed region of $(\epsilon_\mu - \epsilon_e, \epsilon_\tau - \epsilon_e)$ is shown. τ, μ denote the constraints obtained from τ, μ leptonic decays. W denotes the bound obtained from $W \rightarrow l_i \bar{\nu}$ decays. The invisible decay width constraints Eq.(26) is also shown for $\epsilon_e = 0$. The dotted line corresponds to the prediction of class B model (NH) case. (See section IV in text.)

the branching fractions of $\tau \rightarrow \mu(e)\nu\bar{\nu}$ and $\mu \rightarrow e\nu\bar{\nu}$ decays. The constraints obtained from τ and μ leptonic decays (W leptonic decays) are summarized in Eq.(13) (Eq.(6)) of [28],

$$\begin{aligned}\epsilon_\mu - \epsilon_e &= 0.0002 \pm 0.0042 \quad (0.002 \pm 0.022), \\ \epsilon_\tau - \epsilon_e &= -0.0008 \pm 0.0044 \quad (-0.058 \pm 0.028).\end{aligned}\tag{25}$$

The off-diagonal elements of the correlation matrix of $\epsilon_\mu - \epsilon_e$ and $\epsilon_\tau - \epsilon_e$ are 0.51 and 0.44, respectively [28]. In Fig.(1), we show the constraints Eq.(25) on the plane $(\epsilon_\mu - \epsilon_e, \epsilon_\tau - \epsilon_e)$ by taking into account the correlations. In the figure, we also show the constraint obtained from Eq.(22) for the case that ϵ_e is vanishing. The constraint of Eq.(22) is written with ϵ_i as

$$(\epsilon_\mu - \epsilon_e) + (\epsilon_\tau - \epsilon_e) = (0.008 \pm 0.004) - 3\epsilon_e.\tag{26}$$

From Fig.1, we can see that the constraints obtained from τ and μ leptonic decays are consistent with Z invisible decay width constraint within 1σ CL under the assumption $\epsilon_e = 0$ while the constraints obtained from W decays are not consistent and $\epsilon_\tau - \epsilon_e < 0$ seems to be required in this case. We come back to the lepton universality constraints when we consider the specific structure of Yukawa-Dirac mass term in the following sections.

The Inami-Lim function $\bar{E}(\frac{M_f^2}{m_W^2})$ is shown in Fig. 2. The typical values for the Inami-Lim




FIG. 2: Inami-Lim functions $-\bar{E}$ (thick solid line) and $10 \times F_2$ (dashed line) as functions of the singlet neutrino mass M_N (GeV).

function are $|\bar{E}| \sim 2.8 \sim 8.1$ for $M = 200 \sim 2000$ (GeV). Finally, from the fact that the factor $u_{hI}u_{II}^*$ depends on the flavor structure of m_D , the following relation

$$2|u_{hI}u_{II}| \leq |u_{hI}|^2 + |u_{II}|^2 \leq \sum_{i=e,\mu,\tau} |u_{iI}|^2 = 1, \quad (27)$$

leads us to the constraint, $|u_{hI}u_{II}|^2 \leq 0.25$. Then, numerical value of $S(h, l)$ for the case with N degenerate M_i and X_i is

$$|S(h, l)|^2 \leq \left(N \frac{X}{M}\right)^2 \bar{E} \left(\frac{M^2}{M_W^2}\right)^2 0.25 \simeq 2 \times 10^{-4} \sim 2 \times 10^{-3}, \quad (28)$$

where we denote $M_i \equiv M (i = 1 \sim N)$ and $X_i \sim X (i = 1 \sim N)$. The upper bound of the branching fractions for $b \rightarrow sl_h^\mp l_l^\pm$ is roughly predicted to be $10^{-11} \sim 10^{-10}$ for $200 \text{ (GeV)} < M < 2000 \text{ (GeV)}$.

III. LEPTONIC FCNC AND QLFV RARE DECAYS WITH LOW MASS SCALE SINGLET MAJORANA NEUTRINOS

We now predict the branching fractions of the QLFV processes more concretely. As previously discussed, the branching fractions can be enhanced for rather large values of X_I/M_I . The large values of X_I/M_I are realized for low scale of M_I , which may not generally be consistent with neutrino data. The large values of X_I/M_I can be consistent with neutrino

	(n_1, n_2, n_3)	$V_0 = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$	flavor dependence of $b \rightarrow sl_h^- l_l^+$
Normal	$(0, \sqrt{\Delta m_{\text{sol}}^2}, \sqrt{\Delta m_{\text{atm}}^2 + \Delta m_{\text{sol}}^2})$		
Class A	$(0, X_{12} , X_3)$	$(\mathbf{u}_1^* \times \mathbf{u}_3^*, \mathbf{u}_1, \mathbf{u}_3) p$	$\mathbf{v}_{h2} \mathbf{v}_{12}^*$
Class B	$(0, X_3, X_{12})$	$(\mathbf{u}_1^* \times \mathbf{u}_3^*, \mathbf{u}_3, \mathbf{u}_1) p$	$\mathbf{v}_{h3} \mathbf{v}_{13}^*$
Inverted	$(\sqrt{\Delta m_{\text{atm}}^2 - \Delta m_{\text{sol}}^2}, \sqrt{\Delta m_{\text{atm}}^2}, 0)$		
Class A	$(X_{12} , X_3, 0)$	$(\mathbf{u}_1, \mathbf{u}_3, \mathbf{u}_1^* \times \mathbf{u}_3^*) p$	$\mathbf{v}_{h1} \mathbf{v}_{11}^*$
Class B	$(X_3, X_{12} , 0)$	$(\mathbf{u}_3, \mathbf{u}_1, \mathbf{u}_1^* \times \mathbf{u}_3^*) p$	$\mathbf{v}_{h2} \mathbf{v}_{12}^*$

TABLE I: The assignment of mass spectrum and MNS matrix.

data when the contributions of X_I to m_{eff} in Eq. (9) are cancelled. Such a cancellation can be achieved by taking two almost degenerate small $M_{1,2}$ and tuning the relative phase between $\mathbf{u}_1$ and $\mathbf{u}_2$ so that those two terms contribute to m_{eff} destructively while keeping X_3 so small that its contribution to m_{eff} is suppressed. Thus, we need some specific flavor structure of Yukawa-Dirac mass term in order to obtain an enhancement of the branching fractions. Let us assume $X_1 \sim X_2 \gg X_3$ so as for $X_{12} \equiv X_1 - X_2$ and X_3 to be of order the light neutrino mass squared differences $\sqrt{\Delta m_{\text{sol}}^2}$ or $\sqrt{\Delta m_{\text{atm}}^2}$. The relative phase of Yukawa-Dirac mass term from two singlet neutrinos N_1 and N_2 is tuned as $\mathbf{u}_2 = i\mathbf{u}_1$. Then, m_{eff} becomes

$$m_{eff} = -(\mathbf{u}_1 \mathbf{u}_1^T X_{12} + \mathbf{u}_3 \mathbf{u}_3^T X_3). \quad (29)$$

We further assume the orthogonality of $\mathbf{u}_1$ and $\mathbf{u}_3$, *i.e.* $\mathbf{u}_1^\dagger \cdot \mathbf{u}_3 = 0$, so that $\mathbf{u}_1$ and $\mathbf{u}_3$ can be directly related to MNS matrix. Then, there exists a massless state due to the alignment of $\mathbf{u}_1$ and $\mathbf{u}_2$, which is assigned to $n_1 = 0$ for normal hierarchy and $n_3 = 0$ for inverted hierarchy. The other two masses are given by X_3 or $|X_{12}|$.

In Table I, we classify the assignment of mass spectrum (n_1, n_2, n_3) and the unitary part of the mixing matrix V_0 , where p is a diagonal Majorana phase which is irrelevant to the QFLV and LFV processes and thus we omit it from now on. Notice that V_0 is identical to the MNS matrix if we neglect its deviation from V_{MNS} , $V_0 - V_{\text{MNS}} = \mathcal{O}(\frac{m_D^2}{M^2})$. In fact, since the QLFV and LFV processes are already in the order of $\frac{m_D^2}{M^2}$ at the leading order, the difference can be safely ignored. The flavor dependence of the amplitudes for the QFLV and LFV processes is then extracted in terms of the mixing angles of the neutrino oscillation

$\mathbf{v}_{\mu 2} \mathbf{v}_{e 2}^*$	$c_{13} s_{\text{sol}} c_{\text{sol}} c_{\text{atm}}$	$0.33 c_{13}$
$\mathbf{v}_{\tau 2} \mathbf{v}_{\mu 2}^*$	$-c_{\text{atm}} s_{\text{atm}} c_{\text{sol}}^2$	-0.35
$\mathbf{v}_{\tau 2} \mathbf{v}_{e 2}$	$-c_{13} s_{\text{atm}} c_{\text{sol}} s_{\text{sol}}$	$-0.33 c_{13}$
$\mathbf{v}_{\mu 3} \mathbf{v}_{e 3}^*$	$c_{13} s_{\text{atm}} s_{13} \exp(i\delta)$	$0.71 s_{13} c_{13} \exp(i\delta)$
$\mathbf{v}_{\tau 3} \mathbf{v}_{\mu 3}^*$	$c_{13}^2 s_{\text{atm}} c_{\text{atm}}$	$0.5 c_{13}^2$
$\mathbf{v}_{\tau 3} \mathbf{v}_{e 3}$	$c_{13} c_{\text{atm}} s_{13} \exp(i\delta)$	$0.71 s_{13} c_{13} \exp(i\delta)$
$\mathbf{v}_{\mu 1} \mathbf{v}_{e 1}^*$	$-c_{13} c_{\text{sol}} s_{\text{sol}} c_{\text{atm}}$	$-0.33 c_{13}$
$\mathbf{v}_{\tau 1} \mathbf{v}_{\mu 1}^*$	$-s_{\text{sol}}^2 c_{\text{atm}} s_{\text{atm}}$	-0.16
$\mathbf{v}_{\tau 1} \mathbf{v}_{e 1}^*$	$c_{13} c_{\text{sol}} s_{\text{sol}} s_{\text{atm}}$	$0.33 c_{13}$

TABLE II: The combinations of $\mathbf{v}_{ij}$ relevant to the flavor dependence of QLFV and LFV decays.

from $V_0 = V_{\text{MNS}}$:

$$V_0 = \begin{pmatrix} c_{13} c_{\text{sol}} & c_{13} s_{\text{sol}} & 0 \\ -s_{\text{sol}} c_{\text{atm}} & c_{\text{atm}} c_{\text{sol}} & c_{13} s_{\text{atm}} \\ s_{\text{sol}} s_{\text{atm}} & -s_{\text{atm}} c_{\text{sol}} & c_{13} c_{\text{atm}} \end{pmatrix} + s_{13} \begin{pmatrix} 0 & 0 & \exp(-i\delta) \\ -c_{\text{sol}} s_{\text{atm}} \exp(i\delta) & -s_{\text{sol}} s_{\text{atm}} \exp(i\delta) & 0 \\ -c_{\text{sol}} c_{\text{atm}} \exp(i\delta) & -s_{\text{sol}} c_{\text{atm}} \exp(i\delta) & 0 \end{pmatrix}, \quad (30)$$

where we take $s_{\text{sol}} = 0.56$, $c_{\text{sol}} = 0.84$ and $s_{\text{atm}} = c_{\text{atm}} = \sqrt{\frac{1}{2}}$.

In Table II, we present the relevant combinations of $\mathbf{v}_{ij}$ which correspond to the flavor dependence shown in the fourth column of Table I. The value of s_{13} is very small and thus we ignore the terms of order $\mathcal{O}(s_{13}^2)$. Then, the suppression factor $S(h, l)$ is approximately given by

$$S(h, l) \simeq \left(\frac{X}{M_1} \bar{E}(x_t, x_c, y_1) + \frac{X}{M_2} \bar{E}(x_t, x_c, y_2) \right) \mathbf{v}_{h\alpha} \mathbf{v}_{l\alpha}^*, \quad (31)$$

where α denotes the index depending on the class and neutrino mass hierarchy, $X_1 \simeq X_2 \equiv X$. And the term proportional to $\frac{X_3}{M_3}$ is not relevant at all and thus ignored. By using Eq. (31), the ratios of the branching fractions are given by

$$\frac{\text{Br}(b \rightarrow s \tau^\pm e^\mp)}{\text{Br}(b \rightarrow s \tau^\pm \mu^\mp)} = \left| \frac{\mathbf{v}_{\tau\alpha} \mathbf{v}_{e\alpha}^*}{\mathbf{v}_{\tau\alpha} \mathbf{v}_{\mu\alpha}^*} \right|^2, \quad (32)$$

$$\frac{\text{Br}(b \rightarrow s \mu^\pm e^\mp)}{\text{Br}(b \rightarrow s \tau^\pm \mu^\mp)} = \frac{P(\frac{m_\mu}{m_b})}{P(\frac{m_\tau}{m_b})} \left| \frac{\mathbf{v}_{\mu\alpha} \mathbf{v}_{e\alpha}^*}{\mathbf{v}_{\tau\alpha} \mathbf{v}_{\mu\alpha}^*} \right|^2, \quad (33)$$

where $P(\frac{m_\mu}{m_b})$, and $P(\frac{m_\tau}{m_b})$ are phase space factors and $\frac{P(\frac{m_\mu}{m_b})}{P(\frac{m_\tau}{m_b})} = 2.74$.

	Class A NH (Class B IH)	Class B NH	Class A IH
$\frac{\text{Br}(b \rightarrow s\tau^\pm e^\mp)}{\text{Br}(b \rightarrow s\tau^\pm \mu^\mp)}$	$\left(\frac{s_{\text{atm}} c_{\text{sol}} s_{\text{sol}}}{c_{\text{atm}} s_{\text{atm}} c_{\text{sol}}^2}\right)^2 = 0.89$	$\left(\frac{c_{\text{atm}} s_{13}}{s_{\text{atm}} c_{\text{atm}}}\right)^2 = 2.0 s_{13}^2$	$\left(\frac{c_{\text{sol}} s_{\text{sol}} s_{\text{atm}}}{s_{\text{sol}}^2 c_{\text{atm}} s_{\text{atm}}}\right)^2 = 4.5$
$\frac{\text{Br}(b \rightarrow s\mu^\pm e^\mp)}{\text{Br}(b \rightarrow s\tau^\pm \mu^\mp)}$	$\left(\frac{s_{\text{sol}} c_{\text{sol}} c_{\text{atm}}}{c_{\text{atm}} s_{\text{atm}} c_{\text{sol}}^2}\right)^2 \frac{P_\mu}{P_\tau} = 2.4$	$\left(\frac{s_{\text{atm}} s_{13}}{s_{\text{atm}} c_{\text{atm}}}\right)^2 \frac{P_\mu}{P_\tau} = 5.5 s_{13}^2$	$\left(\frac{c_{\text{sol}} s_{\text{sol}} c_{\text{atm}}}{s_{\text{sol}}^2 c_{\text{atm}} s_{\text{atm}}}\right)^2 \frac{P_\mu}{P_\tau} = 12.0$

TABLE III: Ratios of the branching fractions of $b \rightarrow sl_h^- l_l^+$. $P_\mu \equiv P(\frac{m_\mu}{m_b})$ and $P_\tau \equiv P(\frac{m_\tau}{m_b})$.

	Class A NH (Class B IH)	Class B NH	Class A IH
$\frac{\text{Br}(\tau \rightarrow e\gamma)}{\text{Br}(\tau \rightarrow \mu\gamma)}$	$\left(\frac{s_{\text{atm}} c_{\text{sol}} s_{\text{sol}}}{c_{\text{atm}} s_{\text{atm}} c_{\text{sol}}^2}\right)^2 = 0.89$	$\left(\frac{c_{\text{atm}} s_{13}}{s_{\text{atm}} c_{\text{atm}}}\right)^2 = 2.0 s_{13}^2$	$\left(\frac{c_{\text{sol}} s_{\text{sol}} s_{\text{atm}}}{s_{\text{sol}}^2 c_{\text{atm}} s_{\text{atm}}}\right)^2 = 4.5$
$\frac{\text{Br}(\mu \rightarrow e\gamma)}{\text{Br}(\tau \rightarrow \mu\gamma)}$	$\left(\frac{s_{\text{sol}} c_{\text{sol}} c_{\text{atm}}}{c_{\text{atm}} s_{\text{atm}} c_{\text{sol}}^2}\right)^2 \frac{\tau_\mu}{\tau_\tau} \left(\frac{m_\mu}{m_\tau}\right)^5 = 5.0$	$\left(\frac{s_{\text{atm}} s_{13}}{s_{\text{atm}} c_{\text{atm}}}\right)^2 \frac{\tau_\mu}{\tau_\tau} \left(\frac{m_\mu}{m_\tau}\right)^5 = 11 s_{13}^2$	$\left(\frac{c_{\text{sol}} s_{\text{sol}} c_{\text{atm}}}{s_{\text{sol}}^2 c_{\text{atm}} s_{\text{atm}}}\right)^2 \frac{\tau_\mu}{\tau_\tau} \left(\frac{m_\mu}{m_\tau}\right)^5 = 25$

TABLE IV: Ratios of the branching fractions of $l_h \rightarrow l_l \gamma$.

In Table III, the numerical results of the ratios of the branching fractions given in Eqs. (32,33) are presented. It can be seen that in Class B model for NH case, only $b \rightarrow s\tau^- \mu^+$ can be much larger than the other channel because of the absence of the suppression factor s_{13}^2 for $\tau\mu$ final states, while the branching fractions of the different channels in models except Class B for NH case are within a factor of 10. Furthermore, as discussed in Ref. [14], there is strong correlation between QLFV processes and LFV radiative decays $l_h \rightarrow l_l \gamma$. Experimentally, there are stringent bounds as $\text{Br}(\mu \rightarrow e\gamma) < 1.2 \times 10^{-11}$ [30], $\text{Br}(\tau \rightarrow \mu\gamma) < 6.8 \times 10^{-8}$ [31] and $\text{Br}(\tau \rightarrow e\gamma) < 1.1 \times 10^{-7} (3.9 \times 10^{-7})$ [32, 33]. The bounds on the LFV processes stringently constrain the branching fractions for QLFV processes.

We note that the bound on $\frac{X}{M}$ from the invisible decay width of Z and the present upper bound on $\mu \rightarrow e\gamma$ are not compatible with each other for Class A and Class B IH case, as shown in Table IV. The branching fraction for $l_h \rightarrow l_l \gamma$ is,

$$\text{Br}(l_h \rightarrow l_l \gamma) = \frac{\alpha^3}{256\pi^2 s_W^4} \frac{m_h^4}{M_W^4} \frac{m_h}{\Gamma_h} |G|^2. \quad (34)$$

Numerically computing the pre-factors, the branching fractions are given by

$$\begin{aligned} \text{Br}(\tau \rightarrow \mu\gamma) &= 5.4 \times 10^{-4} |G(\tau, \mu)|^2, \\ \text{Br}(\tau \rightarrow e\gamma) &= 5.4 \times 10^{-4} |G(\tau, e)|^2, \\ \text{Br}(\mu \rightarrow e\gamma) &= 3.1 \times 10^{-3} |G(\mu, e)|^2, \end{aligned} \quad (35)$$

where G is a suppression factor defined by

$$\begin{aligned} G &= \sum_{\alpha=4}^{3+N} V_{l\alpha} V_{h\alpha}^* F_2(y_\alpha) \\ &= \sum_{I=1}^N u_{lI} u_{hI}^* \frac{X_I}{M_I} F_2(y_I), \end{aligned} \quad (36)$$

where $y_I = \frac{M_I^2}{m_W^2}$ and F_2 is Inami-Lim function, $F_2(y) = -\frac{2y^3+5y^2-y}{4(1-y)^3} - \frac{3y^3 \log y}{2(y-1)^4}$, as shown in Fig. 2.

With $M_1 = M_2 = M \ll M_3$ and $X_1 \simeq X_2 = X \gg X_3$, for Class A and Class B (IH) model, we predict,

$$\text{Br}(\mu \rightarrow e\gamma) = 3.1 \times 10^{-3} \left(\frac{2X}{M} \right)^2 (0.33)^2 F_2(y)^2 \geq 4.4 \times 10^{-10}, \quad (37)$$

where we have used $\frac{2X}{M} \geq 0.004$ and $M \geq 200$ (GeV). Therefore, Class A and Class B (IH) models are excluded. If the bound on $\frac{X}{M}$ in Eq. (22) is not taken into account, one can obtain from Eqs. (17,35),

$$\begin{aligned} \text{Br}(b \rightarrow s\tau^- \mu^+) &= 1.9 \times 10^{-4} \text{Br}(\tau \rightarrow \mu\gamma) \frac{|S(\tau\mu)|^2}{|G(\tau\mu)|^2} \leq 1.3 \times 10^{-11} \left| \frac{S(\tau\mu)}{G(\tau\mu)} \right|^2, \\ \text{Br}(b \rightarrow s\tau^- e^+) &= 1.9 \times 10^{-4} \text{Br}(\tau \rightarrow e\gamma) \frac{|S(\tau e)|^2}{|G(\tau e)|^2} \leq 2.0 \times 10^{-11} \left| \frac{S(\tau e)}{G(\tau e)} \right|^2, \\ \text{Br}(b \rightarrow s\mu^- e^+) &= 9.0 \times 10^{-5} \text{Br}(\mu \rightarrow e\gamma) \frac{|S(\mu e)|^2}{|G(\mu e)|^2} \leq 1.1 \times 10^{-15} \left| \frac{S(\mu e)}{G(\mu e)} \right|^2. \end{aligned} \quad (38)$$

While Eq. (38) depends neither on the mass spectrum of heavy Majorana neutrinos nor on the flavor structure of Yukawa-Dirac mass terms, the ratio of $|\frac{S}{G}|$ depends on the details of them. For the present case with $M_1 = M_2 = M \ll M_3$ and $X_1 = X_2 = X \gg X_3$, the ratio $|S(h, l)/G(h, l)|$ is simply given as,

$$\left| \frac{S(h, l)}{G(h, l)} \right|^2 = \left| \frac{\bar{E}(x_t, x_c, y)}{F_2(y)} \right|^2 = 98 \ (M = 200 \text{ GeV}) \sim 285 \ (M = 2000 \text{ GeV}). \quad (39)$$

Therefore, the upper bounds on the branching fractions given in Eq. (38) are translated to

$$\begin{aligned} \text{Br}(b \rightarrow s\tau^- e^+) &\leq (1.3 \sim 3.6) \times 10^{-9} \\ \text{Br}(b \rightarrow s\tau^- \mu^+) &\leq (2.0 \sim 5.0) \times 10^{-9} \\ \text{Br}(b \rightarrow s\mu^- e^+) &\leq (1.1 \sim 3.0) \times 10^{-13}. \end{aligned} \quad (40)$$

The range of the upper bounds corresponds to $M = 200 \sim 2000$ (GeV). For Class A and Class B with IH case, by combining $\mu \rightarrow e\gamma$ upper limit, we may obtain much tighter bound

on the other LFV and QLFV processes. Let us consider Class A (NH) and Class B (IH) cases. From Table III and Table IV, the upper bounds on $\text{Br}(\mu \rightarrow e\gamma)$ and $\text{Br}(b \rightarrow s\mu^-e^+)$ obtained in Eq. (40) severely constrain the other modes,

$$\begin{aligned}\text{Br}(b \rightarrow s\tau^-e^+) &\leq (0.41 \sim 1.1) \times 10^{-13}, \\ \text{Br}(b \rightarrow s\tau^-\mu^+) &\leq (0.46 \sim 1.25) \times 10^{-13}, \\ \text{Br}(\tau \rightarrow \mu\gamma) &\leq 2.4 \times 10^{-12}, \\ \text{Br}(\tau \rightarrow e\gamma) &\leq 2.1 \times 10^{-12}.\end{aligned}\tag{41}$$

The bounds similar to Eq. (41) are obtained for Class A IH case,

$$\begin{aligned}\text{Br}(b \rightarrow s\tau^-e^+) &\leq (0.41 \sim 1.1) \times 10^{-13}, \\ \text{Br}(b \rightarrow s\tau^-\mu^+) &\leq (0.92 \sim 2.5) \times 10^{-14}, \\ \text{Br}(\tau \rightarrow \mu\gamma) &\leq 4.8 \times 10^{-13}, \\ \text{Br}(\tau \rightarrow e\gamma) &\leq 2.2 \times 10^{-12}.\end{aligned}\tag{42}$$

We note that the upper bound of the branching fractions of QLFV processes for Class A (NH, IH) and Class B (IH) models are $10^{-14} \sim 10^{-13}$ and the branching fractions for LFV processes is $10^{-13} \sim 10^{-12}$. As we have already shown in Eq.(37), the Class A model and Class B model for IH case can not satisfy the upper limit of the branching fraction of $\mu \rightarrow e\gamma$ and constraint from the effective light neutrinos number N_ν simultaneously. This is because the former requires very small $\frac{X}{M}$, while the latter requires larger $\frac{X}{M}$. Below, we show Class B model for NH case may satisfy both constraints. Furthermore, the model predicts the large branching fractions of $\tau \rightarrow \mu\gamma$ and $b \rightarrow s\tau\mu$ which are within the reach of near future Super B factories [34, 35]. If we take into account the constraints coming from $\text{Br}(\mu \rightarrow e\gamma)$ and the effective number of light neutrinos N_ν , we can have the parameter regions consistent with the present bounds only for Class B model with NH case. In this class, the stringent experimental limit on $\text{Br}(\mu \rightarrow e\gamma)$ is not effective on $\text{Br}(\tau \rightarrow \mu\gamma)$ and $\text{Br}(b \rightarrow s\tau\mu)$, because the former process is proportional to s_{13}^2 and thus ignorable, but the latter processes are not suppressed by the factor.

In Fig. 3, we have shown the correlation of branching fractions between $\mu \rightarrow e\gamma$ and the other LFV and QLFV processes in Class B model for NH case. The numerical results in Fig. 3 are obtained as follows: We first set $X_1 \sim X_2 = X$ and $M_2 = M_1 = M$. From the constraint given in Eq. (22), the allowed range of M is ,

$$\frac{m_{D1}}{\sqrt{\sigma_{\max}}}\sqrt{2} < M < \frac{m_{D1}}{\sqrt{\sigma_{\min}}}\sqrt{2},\tag{43}$$




FIG. 3: Correlation between the branching fraction for $\mu \rightarrow e\gamma$ and the branching fractions for $b \rightarrow s\tau\mu$ (thick solid line) $b \rightarrow s\tau e$ (solid line) $b \rightarrow s\mu e$ (thin solid line), $\tau \rightarrow \mu\gamma$ (long dashed line) $\tau \rightarrow e\gamma$ (dashed line) for Class B model with NH case. From left to right, the lines correspond to $s_{13} = 0.02, 0.05, 0.1$, respectively. The shaded region is excluded by the current bound on $Br(\mu \rightarrow e\gamma)$.

with $\sigma_{\min} = 0.004, \sigma_{\max} = 0.012$. When we fix m_{D1} , the allowed range of M is determined. By varying M within the above range, we plot the correlation between $Br(\mu \rightarrow e\gamma)$ and the other five QLFV and LFV branching fractions. Here, s_{13} is a free parameter and is chosen to be 0.02, 0.05 and 0.1. m_{D1} is chosen to be 100 (GeV). As can be seen in Fig. 3, the present upper limit on $Br(\mu \rightarrow e\gamma)$ gives a very tight bound on s_{13} , typically smaller than 0.02. With this small s_{13} , $\tau \rightarrow e\gamma$ and $b \rightarrow s\tau(\mu)e$ are also severely suppressed. Only $b \rightarrow s\tau\mu$ and $\tau \rightarrow \mu\gamma$ are free from the suppression and the former branching fraction can be as large as 10^{-10} and the latter can be 10^{-9} . They are independent on small s_{13} .

We show in Fig. 4 and Fig. 5 the dependence of the branching fractions of $b \rightarrow s\tau\mu$ and $\tau \rightarrow \mu\gamma$ on m_{D1} and the heavy Majorana neutrino mass for the exact degenerate case, *i.e.* $M_1 = M_2 = M$. We fix m_{D1} to 20, 50, 100, 200 (GeV). Although the branching fractions become small as m_{D1} becomes small, the change of the branching fractions is within a factor 10. We also consider the non-degenerate case for Majorana neutrino masses ($M_1 \equiv M \neq M_2$) while keeping the degeneracy $X_1 \sim X_2 = X$. By setting $M_2 \equiv RM_1$ and $M_1 \equiv M$, the dependence of the branching fraction $b \rightarrow s\tau\mu$ on the ratio R is studied. The




FIG. 4: $\text{Br}(b \rightarrow s\tau\mu)$ vs. M for Class B (NH). From left to right, the curves correspond to $m_{D1} = 20, 50, 100, 200$ (GeV), respectively.




FIG. 5: $\text{Br}(\tau \rightarrow \mu\gamma)$ vs. M for Class B (NH). From left to right, the curves correspond to $m_{D1} = 20, 50, 100, 200$ (GeV), respectively.

allowed range of the lightest heavy Majorana neutrino mass M of Eq. (43) is modified as $\frac{m_{D1}}{\sqrt{\sigma_{\max}}} \sqrt{1 + \frac{1}{R}} < M < \frac{m_{D1}}{\sqrt{\sigma_{\min}}} \sqrt{1 + \frac{1}{R}}$. From Fig. 6, we find that the lower and the upper limits of M become smaller as R becomes larger. However, the branching fraction does not change so significantly.

IV. SUMMARY AND DISCUSSION

As shown, the contributions of the singlet Majorana neutrinos to QLFV and LFV decays can be significant in the low scale seesaw model motivated by resonant leptogenesis. The




FIG. 6: $\text{Br}(b \rightarrow s\tau\mu)$ vs. M for Class B (NH). We fix $m_{D1} = 100$ (GeV). From right to left, the curves correspond to the ratio $R = \frac{M_2}{M}$ 1, 2, 10, respectively.

branching fractions of inclusive decays $b \rightarrow sl_h^\pm l_l^\mp$ in the seesaw model considered in this paper depends on the suppression factor $\frac{m_D}{M}$ which is arisen from the mixing between the singlet heavy neutrinos with three light neutrinos, and can be as large as about 10% without being in conflict with the neutrino mass squared differences from neutrino data and the current bound on the invisible decay width of Z boson.

We have classified four classes of the model along with the light neutrino mass spectrum and the assignment of the mixing matrix V_0 , and studied how the ratios of the branching fractions for the various channels of QLFV and LFV decays along with lepton flavors could be distinctively predicted in each class. We have found that only the class B for NH case presented in Table I survives the current limit on $\text{Br}(\mu \rightarrow e\gamma)$ and the invisible decay width of Z boson. One may check if the model is consistent with the experiments of lepton universality tests. The class B for NH case predicts

$$\begin{aligned}\epsilon_e &= \frac{2X}{M} s_{13}^2 \leq 5 \times 10^{-6} \quad (s_{13} \leq 0.02), \\ \epsilon_\mu &= \frac{2X}{M} s_{\text{atm}}^2 c_{13}^2 \simeq 0.004, \\ \epsilon_\tau &= \frac{2X}{M} c_{\text{atm}}^2 c_{13}^2 \simeq 0.004.\end{aligned}\tag{44}$$

The model predicts very small ϵ_e and $\epsilon_\mu = \epsilon_\tau$ which are shown in Fig.(1) with dotted line. The model is consistent with the constraint of Z invisible decay width and the lepton universality constraints from τ and μ decays while it may not be consistent with the lepton universality constraints determined by W decays. In this class, the branching fractions of $b \rightarrow s\tau\mu$ and $\tau \rightarrow \mu\gamma$ are predicted to be as large as 10^{-10} and 10^{-9} , respectively. Such large

branching fractions can be tested in the future B factory experiments. The enhancement of the branching fractions of QFLV and LFV is originated from the one loop Feynman diagrams in which the heavy Majorana neutrinos contribute to and it is non-susy contribution.

Acknowledgements

We thank K. Homma and K. Ishikawa for useful discussions. C.S.K. was supported in part by JSPS, in part by CHEP-SRC Program, in part by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) No. R02-2003-000-10050-0 and in part by No. KRF-2005-070-C00030. T.M. is supported by the kakenhi of MEXT, Japan, No. 16028213. S.K.K. was supported in part by BK21 program of the Ministry of Education in Korea.

-
- [1] P. Minkowski, Phys. Lett. **B67**, 421 (1977).
 - [2] T. Yanagida, in the proceedings of the Workshop on Unified Theories and Baryon Number in the Universe, edited by O. Sawada and A. Sugamoto, 95 (1979).
 - [3] M. Gell-Mann, P. Ramond and R. Slansky, in Supergravity, P. van Nieuwenhuizen and D.Z. Freedman (eds.), North Holland Publ. Co.,(1979).
 - [4] R. N. Mohapatra and G. Senjanovich, Phys. Rev. Lett. **44**, 912 (1980).
 - [5] M. Fukugita and T. Yanagida, Phys. Lett. **B174**, 45 (1986).
 - [6] A. Pilaftsis, Int. J. Mod. Phys. **A14**, 1811 (1999).
 - [7] A. Pilaftsis and T. E. J. Underwood, Nucl. Phys. **B692**, 303 (2004).
 - [8] R. Gonzalez Felipe, F. R. Joaquim, and B. M. Nobre, Phys. Rev. **D70**, 085009 (2004).
 - [9] G.C. Branco, R. Gonzalez Felipe, F. R. Joaquim and B. M. Nobre, [Archive:hep-ph/0507092].
 - [10] M. Raidal, A. Strumia, and K. Turzyski, Phys. Lett. **B609**, 351 (2005), [Archive:hep-ph/0408015].
 - [11] S. M. West, Phys. Rev. **D71**, 013004 (2005), [Archive: hep-ph/0408318].
 - [12] T. Hambye, J. March-Russell, S. M. West, JHEP **0407**, 070 (2004), [Archive:hep-ph/0403183].
 - [13] J. March-Russell, S. M. West, Phys. Lett. **B593**, 181 (2004), [Archive:hep-ph/0403067].
 - [14] Xiao-Gang He, G. Valencia and Yili Wang, Phys. Rev. **D70**, 113011 (2004).
 - [15] Belle Collaboration: A. Ishikawa *et al.*, Phys. Rev. Lett. **91**, 261601 (2003).

- [16] B. W. Lee, S. Pakvasa, R.E. Shrock and H. Sugawara, Phys. Rev. Lett. **38**, 937 (1977).
- [17] T. Inami and C. S. Lim, Prog. Theor. Phys. **67**, 1569 (1982).
- [18] G. Cvetič, C. O. Dib, C. S. Kim and J. D. Kim, Phys. Rev. D **71**, 113013 (2005) [arXiv:hep-ph/0504126]; G. Cvetič, C. Dib, C. S. Kim and J. D. Kim, Phys. Rev. D **66**, 034008 (2002) [Erratum-ibid. D **68**, 059901 (2003)] [arXiv:hep-ph/0202212].
- [19] A. Pilaftsis, Phys. Rev. Lett. **95**, 081602 (2005).
- [20] A. Pilaftsis and T.E.J. Underwood, [arXiv:hep-ph/0506107].
- [21] G. C. Branco, T. Morozumi, B. Nobre and M. N. Rebelo, Nucl. Phys. **B617**, 475 (2001).
- [22] A. Broncano, M. B. Gavela and E. Jenkins, Nucl. Phys. **B672**, 163 (2003).
- [23] T. Fujihara, S. Kaneko, S. Kang, D. Kimura, T. Morozumi and M. Tanimoto, Phys. Rev. D **72**, 016006 (2005) [arXiv:hep-ph/0505076].
- [24] T. Inami and C. S. Lim, Prog. Theor. Phys. **65**, 297 (1981); **65**, 1772(E) (1981).
- [25] Z. Gagy-Palfy, A. Pilaftsis and K. Schilcher, Phys. Lett. **B343**, 275 (1995).
- [26] Will Loinaz, N. Okamura, S. Rayyan and T. Takeuchi, Phys. Rev. **D68**, 073001 (2003).
- [27] S. L. Glashow, [arXiv:hep-ph/0301250].
- [28] Will Loinaz, N. Okamura, S. Rayyan and T. Takeuchi Phys. Rev. **D70**, 113004 (2004).
- [29] The LEP Collaborations ALEPH, DELPHI, L3, OPAL, the LEP Electroweak Working Group, the SLD Electroweak and Heavy Flavor Groups, [arXiv:hep-ex/0312023].
- [30] MEGA Collaboration: M. L. Brooks *et al.*, Phys. Rev. Lett. **83**, 1521 (1999); MEGA Collaboration: M. Ahmed *et al.*, Phys. Rev. **D65**, 112002 (2002).
- [31] BABAR Collaboration: B. Aubert *et al.*, Phys. Rev. Lett. **95**, 041802 (2005).
- [32] BABAR Collaboration: B. Aubert *et al.*, [arXiv:hep-ex/0508012].
- [33] Belle Collaboration: K. Hayasaka *et al.*, Phys. Lett. **B613**, 20 (2005).
- [34] Physics at Super *B* Factory, by SuperKEKB Physics Working Group, A. G. Akeroyd *et al.*, [arXiv:hep-ex/0406071].
- [35] The Discovery Potential of a Super *B* Factory. Proceedings, SLAC Workshops, Stanford, USA (2003), J. Hewett, (ed.) *et al.*, [arXiv:hep-ph/0503261].